Iron (Native Iron) - Fe

Hardness: 4.5. Cleavage: Imperfect/fair. Insoluble in water. Rapidly dissolves in acids, especially HNO₃. Slowly soluble in acetic acid. Insoluble in concentrated H₂SO₄ but this treatment is said to be "quite unthinkable." Insoluble in alkalis, ethanol or acetone. May oxidize (rust) in moist atmosphere; store in sealed container over desiccant. Avoid wet cleaning methods where possible. "A bright dip for iron minerals" consists of two ounces oxalic acid, 100 mL 30% H₂O₂ (12% H₂O₂ much safer and likely equally effective), 875 mL room temperature water and 25 mL H₂SO₄ (concentration unspecified but probably concentrated H₂SO₄). "Iron of archaeological value can be cleaned of rust by contact with granulated Zn in the presence of (concentration unspecified) NaOH." Alternately, "Iron artifacts may be cleaned chemically by several methods. The objects should first be freed of dirt and loose rust [with] brushes and steel wool. For small objects, or a small quantity, rust may be removed [with] a solution of two parts SnCl₂ to eight parts of water."

References

Mindat: https://www.mindat.org/min-2047.html

Handbook of Mineralogy: https://www.handbookofmineralogy.org/pdfs/iron.pdf

Burns, N. (1941) National Park Service Field Manual for Museums, 426 p. United States Government Printing Office, Washington, D. C.

Hardinger, S. (2025) Mineral Specimen Cleaning and Development for the Amateur, 339 p.

Pearl, R. (1972) Cleaning and Preserving Minerals Fifth Revised Edition, 87 p. Earth Science Publishing Company, Colorado Springs, Colorado.

Scott, A. (1926) The Cleaning and Restoration of Museum Exhibits, 60 p. His Majesty's Stationery Office, London.

Sinkankas, J. (1972) Gemstone & Mineral Data Book, 346 p. Winchester Press, New York.

Weast, R., Ed. (1982) CRC Handbook of Chemistry and Physics, 2380 p., CRC Press, Inc., Cleveland.